



Calculated Barrier Heights for OH + C₂H₂ and OH + C₂H₄ Using Unrestricted Møller–Plesset Perturbation Theory with Spin Annihilation

Carlos Sosa[†] and H. Bernhard Schlegel^{*‡}

Contribution from the Department of Chemistry, Wayne State University, Detroit, Michigan 48202. Received October 6, 1986

Abstract: The reactions of OH radical with C₂H₂ and C₂H₄ have been studied with ab initio molecular orbital techniques. Reactants, loose clusters, transition structures, and products were optimized at UHF/3-21G and UHF/6-31G*. The barrier heights have been computed by using unrestricted Hartree–Fock and Møller–Plesset perturbation theory up to fourth order, including single, double, and quadruple excitations. Spin contamination in the UHF wave function has been corrected by annihilating the largest spin contaminant. The vibrational frequencies were computed by using analytical derivative methods at the UHF/3-21G level. The barrier heights for both reactions are overestimated by 7–15 kcal/mol at the UMP2, UMP3, and UMP4 levels. Annihilation of the largest spin contaminant lowers the barrier heights by 7–15 kcal/mol. Calculations at the PMP4/6-31G* level are in good agreement with the estimated experiment barrier heights.

The reactions of OH with acetylene and ethylene are known to be important in hydrocarbon combustion as well as atmospheric chemistry.^{1–5} Experimental studies on OH + C₂H₂^{1–12} and OH + C₂H₄^{1–4,11–25} have shown that near room temperature the predominant mechanism involves the electrophilic addition of the OH radical to the π bond, forming an activated complex which can be collisionally stabilized.



In early experimental work on OH + C₂H₂, no pressure dependence of the rate constant was found at low temperatures.^{6,7} However, recent studies covering a wider range of temperature and pressure, and using flash photolysis/resonance fluorescence and laser pyrolysis/laser fluorescence techniques,^{8–10} indicate that the rate constant does depend on the pressure, consistent with eq 1 and 2. Analysis of the temperature and pressure dependence yields an activation energy of 1.3 ± 0.1 kcal/mol,^{9,10} based on an estimated heat of reaction of -36 ± 6 kcal/mol for OH + C₂H₂ → C₂H₂OH. At temperatures higher than 1000 K, the addition channel becomes less important because of competition from hydrogen abstraction (estimated activation energy 6–8 kcal/mol).^{9,10}

The OH + C₂H₄ system has also received considerable attention.^{11–25} In accord with eq 1 and 2, the rate for OH + C₂H₄ is found to be dependent on the total pressure. However, unlike addition to acetylene, a small negative activation energy has been found (-0.9 ± 0.2,^{13,14} -0.7 ± 0.3,¹⁵ -0.6 ± 0.3¹⁶ kcal/mol). Several explanations have been given for this, including the formation of a weakly bound complex.²⁶ Estimates of the heat of reaction for the addition process range from -29 to -32 kcal/mol.^{4,16–18} Mass spectral methods have been used to observe directly the primary adduct, C₂H₄OH, and to study its decomposition into CH₃ + CH₂O and H + CH₃CHO.^{18,19} At higher temperatures, hydrogen abstraction becomes the dominant pathway (estimated activation energy ca. 3 kcal/mol).^{1,24,25}

In an earlier theoretical study, Melius, Binkley, and Koszykowski²⁷ examined the reactions of OH with C₂H₂, C₂H₄, and HCN. Geometries were optimized at the HF/6-31G* level and energies were calculated by using fourth order unrestricted Møller–Plesset perturbation theory (UMP4/6-31G**). The heats of reaction and barrier heights were estimated by applying bond additivity corrections to the UMP4 calculations (BAC–MP4). These corrections were determined by a least-squares fit to ca. 50 molecules with well-established heats of formation. For

open-shell molecules, estimated corrections for contamination from higher spin states are also included.

In previous work we have shown that the barrier heights for hydrogen addition to ethylene and formaldehyde are overestimated by 6–10 kcal/mol at the UMP4 level due to spin contamination in the UHF wave function. The difficulties in computing barrier heights with unrestricted Møller–Plesset perturbation theory can be overcome by annihilation of the largest spin contaminant. This leads to a much improved description of bond dissociation curves²⁸

(1) *Combustion Chemistry*; Gardiner, W. C., Jr., Ed.; Springer-Verlag: New York, 1984.

(2) Hucknall, D. J. *Chemistry of Hydrocarbon Combustion*; Chapman and Hall: New York, 1985.

(3) Glassman, I. *Combustion*; Academic Press: New York, 1977.

(4) Kerr, J. A.; Parsonage, M. J. *Evaluated Kinetic Data on Gas Phase Addition Reactions*; Butterworths: England, 1972.

(5) Williams, A.; Smith, D. B. *Chem. Rev.* **1970**, *70*, 267.

(6) Breen, J. E.; Glass, G. P. *Int. J. Chem. Kinet.* **1970**, *3*, 145.

(7) Pastrana, A.; Carr, R. W., Jr. *Int. J. Chem. Kinet.* **1974**, *6*, 587.

(8) Perry, R. A.; Atkinson, R. A.; Pitts, J. N., Jr. *J. Chem. Phys.* **1977**, *67*, 5577.

(9) Michael, J. V.; Nava, D. F.; Borkowski, R. P.; Payne, W. A.; Stief, I. J. *J. Chem. Phys.* **1980**, *73*, 6108.

(10) Smith, G. P.; Fairchild, P. W.; Crosley, D. R. *J. Chem. Phys.* **1984**, *81*, 2667.

(11) Smith, I. W. M.; Zellner, R. *J. Chem. Soc., Faraday Trans. 2* **1973**, *69*, 1617.

(12) Davis, D. D.; Fischer, S.; Schiff, R.; Watson, R. T.; Bollinger, W. J. *Chem. Phys.* **1975**, *63*, 1707.

(13) Greiner, N. R. *J. Chem. Phys.* **1970**, *53*, 1284.

(14) Tully, F. P. *Chem. Phys. Lett.* **1983**, *96*, 148.

(15) Atkinson, R.; Perry, R. A.; Pitts, J. N., Jr. *J. Chem. Phys.* **1977**, *66*, 1197.

(16) Zellner, R.; Lorenz, K. *J. Phys. Chem.* **1984**, *88*, 984.

(17) Howard, C. J. *J. Chem. Phys.* **1976**, *65*, 4771.

(18) Bartels, M.; Hoyerman, K.; Sievert, R. *Symp. (Int.) Combust. [Proc.]* **1982**, *19*, 61.

(19) Morris, E. D., Jr.; Stedman, D. H.; Niki, H. *J. Am. Chem. Soc.* **1971**, *93*, 3570.

(20) Morris, E. D., Jr.; Niki, H. *J. Phys. Chem.* **1971**, *75*, 3640.

(21) Lloyd, A. C.; Darnall, K. R.; Winer, A. M.; Pitts, J. N., Jr. *J. Phys. Chem.* **1976**, *80*, 789.

(22) Atkinson, R.; Darnall, K. R.; Lloyd, A. C.; Winer, A. M.; Pitts, J. N., Jr. *Adv. Photochem.* **1979**, *11*, 375.

(23) Klein, Th.; Barnes, I.; Becker, K. H.; Fink, E. H.; Zabel, F. *J. Phys. Chem.* **1984**, *88*, 5020.

(24) Bradley, J. N.; Capey, W. D.; Fair, R. W.; Pritchard, D. K. *Int. J. Chem. Kinet.* **1976**, *8*, 549.

(25) Westbrook, C. K.; Dryer, F. L.; Schug, K. P. *Symp. (Int.) Combust. [Proc.]* **1982**, *19*, 153.

(26) Singleton, D. L.; Cvetonovic, R. J. *J. Am. Chem. Soc.* **1976**, *98*, 6812.

(27) Melius, C. F.; Binkley, J. S.; Koszykowski, M. L. *8th International Symposium on Gas Kinetics*; Nottingham: England, 1984, and unpublished results.

[†] Present address: Quantum Theory Project, University of Florida, Gainesville, Florida 32605.

[‡] Camille and Henry Dreyfus Teacher-Scholar.

and much better agreement between experimental and theoretical barrier heights.²⁹ The present study uses these same techniques to investigate the structures and energetics of the reactants, loose clusters, transition states, and radical intermediates for the reactions $\text{OH} + \text{C}_2\text{H}_2 \rightarrow \text{C}_2\text{H}_2\text{OH}$ and $\text{OH} + \text{C}_2\text{H}_4 \rightarrow \text{C}_2\text{H}_4\text{OH}$.

Method

Ab initio molecular orbital calculations were performed with the GAUSSIAN 82 system of programs.³⁰ The restricted Hartree-Fock method (RHF) was used for closed shell systems and the unrestricted Hartree-Fock method (UHF) for open shell systems.³¹ All equilibrium geometries and transition structures were fully optimized at the Hartree-Fock level by using analytical gradient methods³² with split-valence (3-21G)³³ and split-valence plus polarization (6-31G*)³⁴ basis sets. In addition, several points were calculated along the reaction paths for the OH addition by fixing the hydroxy-reactant distance, $R(\text{C}-\text{O})$, and minimizing the energy with respect to all other parameters. Vibrational frequencies and zero-point energies were obtained from analytical second derivatives³⁵ calculated at the HF/3-21G level. Electron correlation energy was calculated by using fourth order Møller-Plesset perturbation theory in the space of single, double, and quadruple excitations (MP4SDQ, frozen core).

The effects of spin contamination were examined by annihilation of the next highest spin component in the UHF wave function. In terms of the Löwdin spin projection operator³⁶

$$\hat{P}_s = \prod_{k \neq s} \frac{\hat{S}^2 - k(k+1)}{s(s+1) - k(k+1)} \quad (3)$$

the projected Hartree-Fock energy can be written as

$$E_{\text{proj HHF}} = \frac{\langle \hat{P}_s \Psi_0 | \hat{H} | \hat{P}_s \Psi_0 \rangle}{\langle \hat{P}_s \Psi_0 | \hat{P}_s \Psi_0 \rangle} \quad (4)$$

Since \hat{S}^2 commutes with \hat{H} and \hat{P}_s is idempotent, the projected Hartree-Fock energy can also be written as

$$\begin{aligned} E_{\text{proj UHF}} &= \frac{\langle \Psi_0 | \hat{H} \hat{P}_s | \Psi_0 \rangle}{\langle \Psi_0 | \hat{P}_s | \Psi_0 \rangle} \\ &= \langle \Psi_0 | \hat{H} | \Psi_0 \rangle + \frac{\sum_{i \neq 0} \langle \Psi_0 | \hat{H} | \psi_i \rangle \langle \psi_i | \hat{P}_s | \Psi_0 \rangle}{\langle \Psi_0 | \hat{P}_s | \Psi_0 \rangle} \end{aligned} \quad (5)$$

Because the UHF wave function satisfies Brillouin's theorem and H contains only 1 and 2 operators, the summation over ψ_i can be restricted to all double excitations.

Often, the largest contribution to the spin contamination comes from only the next highest spin, i.e., $s+1$. Under such circumstances, the full projection operator can be approximated by the first term in eq 3, $k = s+1$. The result is no longer a projector (not idempotent) but an annihilation operator, \hat{A}_{s+1} , that removes the $s+1$ spin contaminant (the denominator is chosen to ensure intermediate normalization of $\hat{A}_{s+1} \Psi_0$).

$$\hat{A}_{s+1} = \frac{\hat{S}^2 - (s+1)(s+2)}{\langle \Psi_0 | \hat{S}^2 | \Psi_0 \rangle - (s+1)(s+2)} \quad (6)$$

(28) Schlegel, H. B. *J. Chem. Phys.* **1986**, *84*, 4530.
 (29) Sosa, C.; Schlegel, H. B. *Int. J. Quantum Chem.* **1986**, *29*, 1001.
 (30) Binkley, J. S.; Frisch, M. J.; DeFrees, D. J.; Raghavachari, K.; Whiteside, R. A.; Schlegel, H. B.; Fluder, E. M.; Pople, J. A. GAUSSIAN 82; Carnegie-Mellon University: Pittsburgh, 1983.

(31) Pople, J. A.; Nesbet, R. K. *J. Chem. Phys.* **1954**, *27*, 571.

(32) Schlegel, H. B. *J. Comput. Chem.* **1982**, *3*, 214.

(33) Binkley, J. S.; Pople, J. A.; Hehre, W. J. *J. Am. Chem. Soc.* **1980**, *102*, 939.

(34) Hehre, W. J.; Ditchfield, R.; Pople, J. A. *J. Chem. Phys.* **1972**, *56*, 2257. Hariharan, P. C.; Pople, J. A. *Chem. Phys. Lett.* **1972**, *66*, 217.

(35) Pople, J. A.; Krishnan, R.; Schlegel, H. B.; Binkley, J. S. *Int. Quantum Chem., Quantum Chem. Symp.* **1979**, *13*, 225.

(36) Löwdin, P.-O. *Phys. Rev.* **1955**, *97*, 1509.




Figure 1. Geometries of the 2B_2 100se clusters: HF/3-21G optimized (no superscript), HF/6-31G* optimized (asterisk), in Å and deg.

This simplification was used in our first two papers,^{28,29} and on the basis of eq 5 it yields the following formulae for the projected Hartree-Fock energy and wave function.

$$\begin{aligned} E_{\text{PUHF}} &= \langle \Psi_0 | \hat{H} | \Psi_0 \rangle + \frac{\sum_{i \neq 0} \langle \Psi_0 | \hat{H} | \psi_i \rangle \langle \psi_i | \hat{A}_{s+1} | \Psi_0 \rangle}{\langle \Psi_0 | \hat{A}_{s+1} | \Psi_0 \rangle} \\ &= \langle \Psi_0 | \hat{H} | \Psi_0 \rangle + \frac{\sum_{i \neq 0} \langle \Psi_0 | \hat{H} | \psi_i \rangle \langle \psi_i | \hat{S}^2 | \Psi_0 \rangle}{\langle \Psi_0 | \hat{S}^2 | \Psi_0 \rangle - (s+1)(s+2)} \\ &= E_{\text{UHF}} + \Delta E_{\text{PUHF}} \end{aligned} \quad (7)$$

$$\hat{A}_{s+1} \Psi_0 = \Psi_0 + \frac{\sum_{i \neq 0} \psi_j \langle \psi_j | \hat{S}^2 | \Psi_0 \rangle}{\langle \Psi_0 | \hat{S}^2 | \Psi_0 \rangle - (s+1)(s+2)} = \Psi_0 + \tilde{\Psi}_1 \quad (8)$$

For $\tilde{\Psi}_1$, ψ_j runs over all single excitations and all $\alpha\beta$ -type double excitations.

Perturbation corrections for electron correlation, Ψ_1 , Ψ_2 , etc., consist of single, double, and higher excitations. As a first approximation to spin-projected UMPn energies, the spin correction $\tilde{\Psi}_1$ must be reduced by the amount already contained in Ψ_1 , Ψ_2 , etc. This leads to the following approximate formulae for the projected MPn energies.

$$E_{\text{PMP2}} = E_{\text{UMP2}} + \Delta E_{\text{PUHF}} \left\{ 1 - \frac{\langle \tilde{\Psi}_1 | \Psi_1 \rangle}{\langle \tilde{\Psi}_1 | \tilde{\Psi}_1 \rangle} \right\} \quad (9)$$

$$E_{\text{PMP3}} = E_{\text{UMP3}} + \Delta E_{\text{PUHF}} \left\{ 1 - \frac{\langle \tilde{\Psi}_1 | \Psi_1 + \Psi_2 \rangle}{\langle \tilde{\Psi}_1 | \tilde{\Psi}_1 \rangle} \right\} \quad (10)$$

$$\begin{aligned} E_{\text{PMP4}} &= E_{\text{UMP4}} + \Delta E_{\text{PUHF}} \left\{ 1 - \frac{\langle \tilde{\Psi}_1 | \Psi_1 + \Psi_2 + \Psi_3 \rangle}{\langle \tilde{\Psi}_1 | \tilde{\Psi}_1 \rangle} \right\} \\ &\approx E_{\text{UMP4}} + \Delta E_{\text{PUHF}} \left\{ 1 - \frac{\langle \tilde{\Psi}_1 | \Psi_1 + \Psi_2 \rangle}{\langle \tilde{\Psi}_1 | \tilde{\Psi}_1 \rangle} \right\} \end{aligned} \quad (11)$$




Figure 2. Geometries of the OH + C₂H₂ and OH + C₂H₄ transition states: HF/3-21G optimized (no superscript), HF/6-31G* optimized (asterisk), in Å and deg.





Figure 3. Geometries of 2-hydroxyethyl and 2-hydroxyethynyl radicals: HF/3-21G optimized (no superscript), HF/6-31G* optimized (asterisk), in Å and deg.

Results and Discussion

Geometry. The optimized geometries at the Hartree-Fock level for the loose clusters, transition structures, and radical intermediates are collected in Figures 1-3. Preliminary calculations on these two systems showed that the minimum energy path is one in which the OH radical is located in the symmetry plane that passes through both carbon atoms in ethylene or acetylene. Within C_s symmetry, OH can approach these molecules with the hydrogen syn or anti to the CC bond. The anti approach leads to a local maxima with two imaginary frequencies, while the syn approach corresponds to the saddle point with only one imaginary frequency. Relative to the reactants, the HF/6-31G* optimized values for




Figure 4. Energy profiles along the reaction path for addition of OH + C₂H₂ computed with the 3-21G basis set.




Figure 5. Energy profiles along the reaction path for addition of OH + C₂H₂ computed with the 6-31G* basis set.




Figure 6. Energy profiles along the reaction path for addition of OH + C₂H₄ computed with the 3-21G basis set.

R(CC), *R*(CH), and *R*(OH) in the transition structure differ by only 5%, 0.3%, and -0.4%, respectively, for OH + C₂H₂ and 6%, -0.3%, and -0.6% for OH + C₂H₄. This small perturbation of the geometrical parameters at the saddle point relative to the reactants suggests that the transition structures at the Hartree-Fock level are located relatively early in the entrance channel.

Table I. Vibrational Frequencies^{a,b}


structures	frequencies (cm ⁻¹)
	Minima
OH	3609 (3735)
C ₂ H ₂	902 (612), 918 (729), 2234 (1974)
	3596 (3282), 3719 (3373)
C ₂ H ₄	944 (826), 1115 (949), 1157 (943),
	1165 (1023), 1387 (1220),
	1522 (1342), 1640 (1444),
	1842 (1630), 3238 (3026),
	3305 (3021), 3371 (3103),
	3404 (3105)
C ₂ H ₂ OH	432, 517, 706, 903, 925, 1115
	1363, 1451, 1631, 3347, 3460, 3890
C ₂ H ₄ OH	215, 372, 435, 525, 908, 993
	1117, 1195, 1293, 1488, 1525, 1576
	1675, 3182, 3219, 3300, 3412, 3862
	Transition Structures
OH + C ₂ H ₂ → HOCHCH	5981, 165, 293, 413, 636, 786
	877, 974, 1682, 3545, 3599, 3697
OH + C ₂ H ₄ → HOCH ₂ CH ₂	556i, 97, 304, 474, 820, 895
	982, 1042, 1091, 1271, 1361, 1620
	1673, 3303, 3337, 3388, 3426, 3690
	Loose Clusters
OH + C ₂ H ₂ → OH...C ₂ H ₂	66, 100, 272, 293, 901, 915
	917, 925, 2224, 3587, 3616, 3710
OH + C ₂ H ₄ → OH...C ₂ H ₄	71, 107, 108, 298, 363, 942
	1131, 1169, 1171, 1388, 1517, 1639
	1829, 3307, 3326, 3379, 3401, 3606

^a Experimental values in parentheses from ref 43. ^b The zero-point energies in (kcal/mol) at the HF/3-21G level are the following: OH (5.2), C₂H₂ (18.9), C₂H₄ (34.6), C₂H₂OH (28.2), C₂H₄OH (43.3), OH + C₂H₂ transition structure (23.8), OH + C₂H₄ transition structure (41.1), OH + C₂H₂ cluster (25.1), and OH + C₂H₄ cluster (41.1).

In order to study the effect of electron correlation and spin contamination on the position of the transition structures, several points along the reaction path were optimized at the Hartree-Fock level, followed by single point calculations at the UMPn and PMPn levels. The results are plotted in Figures 4-7. Correlation corrections appear to displace the transition structure by 0.01-0.03 Å toward a later transition state for OH + C₂H₂, but 0.1-0.3 Å toward an earlier transition structure for OH + C₂H₄. For both systems annihilation of the largest spin contaminant shifts the transition structure by about 0.10-0.25 Å toward the reactants.

Two different loose clusters were optimized for both OH + C₂H₂ and OH + C₂H₄ (Figure 1). The binding energies are small and the monomer geometries are essentially unperturbed. For each system the ²B₁ and ²B₂ states are nearly identical in energy and structure, differing only in the orientation of the half-filled π orbital of the hydroxy. Although the ²B₁ states are 0.1-0.3 kcal/mol more stable, the ²B₂ states correlate with the ²A' electronic configurations of the transition structures. The structures of the OH + C₂H₂ and OH + C₂H₄ clusters closely resemble hydrogen fluoride complexes with acetylene and ethylene, which have been studied both experimentally³⁷⁻³⁹ and theoretically.^{40,41}

Several conformations of the radical intermediates were examined. Structures I-IV represent the four different planar conformations for the C₂H₂OH radical. The optimized geometries are very similar to vinyl radical⁴² and vinyl alcohol.⁴³ Structures II and IV are local maxima with imaginary frequencies of 65i and 47i cm⁻¹ corresponding to OH torsion. Structures I and III

(37) Read, W. G.; Flygare, W. H. *J. Chem. Phys.* **1982**, *76*, 2238.(38) Shea, J. A.; Flygare, W. H. *J. Chem. Phys.* **1982**, *76*, 4857.(39) Nelson, D. D., Jr.; Fraser, G. T.; Klempner, W. J. *J. Chem. Phys.* **1985**, *82*, 4483.(40) Pople, J. A.; Frisch, M. J.; Del Bene, J. E. *J. Chem. Phys. Lett.* **1982**, *91*, 185.(41) Frisch, M. J.; Pople, J. A.; Del Bene, J. E. *J. Chem. Phys.* **1983**, *78*, 4063.(42) Dupuis, M.; Wendoloski, J. J. *J. Chem. Phys.* **1984**, *80*, 5696.(43) Stull, D. R.; Prophet, H. *JANAF Thermochemical Tables*; U.S. Government Printing Offices: Washington DC, 1971, and subsequent supplements.**Figure 7.** Energy profiles along the reaction path for addition of OH + C₂H₄ computed with the 6-31G* basis set.**Table II.** Total Energies for Reactants, Clusters, and Products of the OH + C₂H₂ System^a

level	reactants ^b	cluster ^{c,d}	products
Without Spin Projection			
UHF/3-21G	-151.36618	-151.37019	-151.39772
MP2/3-21G	-151.63196	-151.63614	-151.65786
MP3/3-21G	-151.64286	-151.64696	-151.67080
MP4/3-21G	-151.65231	-151.65627	-151.68199
UHF/6-31G*	-152.20011	-152.20409	-152.24291
MP2/6-31G*	-152.58527	-152.59031	-152.63386
MP3/6-31G*	-152.60761	-152.61252	-152.65404
MP4/6-31G*	-152.61454	-152.61925	-152.66272
After Annihilation of the Largest Spin Contaminant			
PUHF/3-21G	-151.36712	-151.37110	-151.41100
PMP2/3-21G	-151.63262	-151.63677	-151.66978
PMP3/3-21G	-151.64330	-151.64736	-151.68044
PMP4/3-21G	-151.65274	-151.65668	-151.69164
PUHF/6-31G*	-152.20275	-152.20671	-152.25499
PMP2/6-31G*	-152.58676	-152.59177	-152.64465
PMP3/6-31G*	-152.60841	-152.61330	-152.66270
PMP4/6-31G*	-152.61535	-152.62002	-152.67138

^a Energy in au. ^b Total energies for reactants with the 6-31G** basis: HF = -152.21016, MP2 = -152.61138, MP3 = -152.63465, and MP4 = -152.64105. ^c ²B₂ state, geometry optimized at UHF/6-31G*. ^d Total energies for the loose cluster with the 6-31G** basis: HF = -152.21403, MP2 = -152.61615, MP3 = -152.63928, and MP4 = -152.64548.

both are local minima with III more stable than I by 1 kcal/mol at the HF/3-21G level. For 2-hydroxyethyl radical, structure V is a local maximum (OH torsion, 311i cm⁻¹); structure VI also corresponds to a maximum (CH₂ torsion, 56i cm⁻¹). Similar to the FCH₂CH₂ radical,⁴⁴ the minimum for HOCH₂CH₂ corresponds to a gauche conformation of the CH₂ group, as indicated in structure VI and Figure 3 (CH₂ torsion 215 cm⁻¹ and OH torsion 372 cm⁻¹).


(44) Bouma, W. J.; Radom, L.; Rodwell, W. R. *Theor. Chim. Acta* **1980**, *56*, 149.

Table III. Total Energies for Reactants, Clusters, and Products of the OH + C₂H₄ System^a

level	reactants ^b	cluster ^{c,d}	products
Without Spin Projection			
UHF/3-21G	-152.571 22	-152.576 61	-152.591 65
MP2/3-21G	-152.835 32	-152.840 55	-152.868 43
MP3/3-21G	-152.858 44	-152.863 46	-152.886 93
MP4/3-21G	-152.867 02	-152.871 86	-152.896 14
UHF/6-31G*	-153.414 00	-153.418 38	-153.443 28
MP2/6-31G*	-153.804 98	-153.810 45	-153.855 68
MP3/6-31G*	-153.837 91	-153.843 12	-153.881 19
MP4/6-31G*	-153.845 09	-153.850 09	-153.888 45
After Annihilation of the Largest Spin Contaminant			
PUHF/3-21G	-152.572 15	-152.577 11	-152.594 36
PMP2/3-21G	-152.835 98	-152.841 18	-152.870 29
PMP3/3-21G	-152.858 87	-152.863 86	-152.888 12
PMP4/3-21G	-152.867 46	-152.872 26	-152.897 33
PUHF/6-31G*	-153.416 64	-153.421 00	-153.446 35
PMP2/6-31G*	-153.806 47	-153.811 90	-153.857 65
PMP3/6-31G*	-153.838 71	-153.843 90	-153.882 38
PMP4/6-31G*	-153.845 89	-153.850 87	-153.889 64

^aEnergy in au. ^bTotal energies for reactants with the 6-31G** basis: HF = -152.883 94, MP2 = -152.848 72, MP3 = -152.883 94, and MP4 = -152.889 99. ^cB₂ state, geometry optimized at UHF/6-31G*. ^dTotal energies for the loose cluster with the 6-31G** basis: HF = -152.431 46, MP2 = -152.853 89, MP3 = -152.888 83, and MP4 = -152.894 67.

Table IV. Heats of Reaction^a

level	OH + C ₂ H ₂	OH + C ₂ H ₄
UHF/3-21G	-19.79	-12.82
MP2/3-21G	-16.25	-20.78
MP3/3-21G	-17.53	-17.88
MP4/3-21G	-18.62	-18.27
PUHF/3-21G	-27.54	-13.94
PMP2/3-21G	-23.32	-21.53
PMP3/3-21G	-23.31	-18.35
PMP4/3-21G	-24.41	-18.74
UHF/6-31G*	-26.86	-18.37
MP2/6-31G*	-30.49	-31.81
MP3/6-31G*	-29.14	-27.16
MP4/6-31G*	-30.23	-27.21
PUHF/6-31G*	-32.78	-18.64
PMP2/6-31G*	-36.33	-32.12
PMP3/6-31G*	-34.07	-27.40
PMP4/6-31G*	-35.16	-27.45
ΔZPE ^b	4.10	3.50
exptl ^c ΔH(0 K)	-34 ± 5	-31 ± 5

^aIn kcal/mol. ^bAt the UHF/3-21G level. ^cReference 10.

Frequencies. The HF/3-21G harmonic frequencies are collected in Table I and are compared with experimental (anharmonic) frequencies,⁴⁵ where available. The computed frequencies are ca. 10–15% too high due to basis set effects and due to neglect of vibrational anharmonicity and electron correlation.^{46,47} The low imaginary frequency of the transition structures for OH adding to acetylene and ethylene indicates a relatively flat potential energy surface for the entrance channel, consistent with Figures 4–7. Because spin projection shifts both transition states to much earlier positions along the reaction coordinate, the actual transition-state frequencies will probably be more reactant like, and the zero point energy corrections to the barriers will be much smaller or negligible.

Heats of Reaction. The total energies for reactants and products are collected in Tables II and III; the heats of reaction are summarized in Table IV. Inspection of the data indicates that

(45) Schlegel, H. B. *J. Phys. Chem.* **1982**, *86*, 4878.

(46) Pople, J. A.; Schlegel, H. B.; Krishnan, R.; DeFrees, D. J.; Binkley, J. S.; Frisch, M. J.; Whiteside, R. A.; Hout, R. F.; Hehre, W. J. *Int. J. Quantum Chem., Quantum Chem. Symp.* **1981**, *15*, 269.

(47) Yamaguchi, Y.; Schaefer, H. F., III *J. Chem. Phys.* **1980**, *73*, 2310.

Table V. Cluster Well Depths^a

level	OH + C ₂ H ₂	OH + C ₂ H ₄
UHF/3-21G	2.52	3.13
MP2/3-21G	2.62	3.28
MP3/3-21G	2.57	3.15
MP4/3-21G	2.48	3.03
UHF/6-31G*	2.50	2.75
MP2/6-31G*	3.16	3.43
MP3/6-31G*	3.08	3.27
MP4/6-31G*	2.96	3.14
UHF/6-31G*	2.43	2.70
MP2/6-31G**	2.99	3.24
MP3/6-31G**	2.91	3.07
MP4/6-31G**	2.78	2.94
PUHF/3-21G	2.50	3.11
PMP2/3-21G	2.60	3.26
PMP3/3-21G	2.55	3.13
PMP4/3-21G	2.47	3.01
PUHF/6-31G*	2.48	2.74
PMP2/6-31G*	3.14	3.41
PMP3/6-31G*	3.07	3.26
PMP4/6-31G*	2.93	3.13
ΔZPE ^b	1.00	1.30

^aB₂ state; energies in kcal/mol. ^bCalculated at the UHF/3-21G level.

Table VI. Relative Energies of OH Addition to C₂H₂ with 3-21G Basis Sets^a

level	R =						
	1.50	1.60	1.80	2.00	2.20	2.50	2.70
Without Spin Projection							
UHF/3-21G	-15.00	-7.47	6.15	8.33	4.15	-0.39	-1.70
MP2/3-21G	-13.22	-6.38	14.14	14.14	5.59	-0.87	-2.34
MP3/3-21G	-14.37	-7.69	10.96	11.80	4.83	-0.87	-2.24
MP4/3-21G	-16.06	-9.91	7.72	9.46	3.78	-1.09	-2.30
After Annihilation of the Largest Spin Contaminant							
PUHF/3-21G	-23.87	-18.51	-15.16	-5.50	-0.70	-1.26	-1.90
PMP2/3-21G	-21.28	-16.34	-5.05	1.90	1.31	-1.68	-2.55
PMP3/3-21G	-20.96	-15.79	-4.73	2.33	1.71	-1.45	-2.39
PMP4/3-21G	-22.66	-18.02	-7.98	-0.01	0.65	-1.68	-2.45

^aEnergies in kcal/mol.

enlarging the basis set (3-21G to 6-31G*) increases the stability of the two radical intermediates (C₂H₂OH and C₂H₄OH) by ca. 5–15 kcal/mol. Electron correlation contributions increase the exothermicity of the reactions by an additional 6 kcal/mol. The heats of reaction predicted at the MP4/6-31G* level are -30.2 kcal/mol for the OH + C₂H₂ system and -27.2 kcal/mol for OH + C₂H₄. Because of greater spin contamination, spin projection stabilized C₂H₂OH (*S*² = 0.992) more than C₂H₄OH (*S*² = 0.763). Changes in zero-point energy contribute 4.1 and 3.5 kcal/mol to ΔH(0 K) for OH + C₂H₂ → C₂H₂OH and OH + C₂H₄ → C₂H₄OH, respectively.

Estimates of the experimental heat of reaction for OH + C₂H₂ → C₂H₂OH include -30,⁸ -36 ± 6,⁹ and -34 ± 5¹⁰ kcal/mol. The present calculations at the PMP4/6-31G* level (corrected for zero-point vibrational energy) predict -31.0 kcal/mol. This agrees well with the experimental estimates and with -33.2 kcal/mol calculated by Melius, Binkley, and Koszykowski²⁷ using the BAC-MP4 approach. For OH + C₂H₄, the experimental estimates for the heat of reaction are -29,¹⁷ -31,¹⁰ and -32 kcal/mol.¹⁴ At the PMP4/6-31G*+ZPE level, the heat of reaction is computed to be significantly less exothermic, -24.0 kcal/mol. This smaller value is supported by the BAC-MP4 calculations²⁷ (-24.9 kcal/mol), which use a quite different approach to correct the energies. However, a more detailed analysis using a larger basis set and relying on isodesmic reactions favors -29 kcal/mol.⁴⁸

The cluster well depths are summarized in Table V. Enlarging the basis set increases the binding energy by only a few tenths

(48) Sosa, C.; Schlegel, H. B. *J. Am. Chem. Soc.*, in press.

Table VII. Relative Energies of OH Addition to C₂H₂ with 6-31G* Basis Sets^a

level	R =						
	1.85	1.95	2.00	2.10	2.30	2.50	2.70
Without Spin Projection							
UHF/6-31G*	10.42	11.99	11.70	9.63	4.73	1.44	-0.35
MP2/6-31G*	13.69	15.98	13.43	8.56	2.59	-0.42	-1.75
MP3/6-31G*	12.24	14.28	12.24	8.23	2.80	-0.16	-1.54
MP4/6-31G*	8.60	11.06	9.56	6.62	2.25	-0.32	-1.57
After Annihilation of the Largest Spin Contaminant							
PUHF/6-31G*	-10.03	-6.23	-2.60	1.94	2.35	0.71	-0.55
PMP2/6-31G*	-5.16	-0.78	0.34	1.44	0.14	-1.36	-2.11
PMP3/6-31G*	-3.22	0.73	1.91	2.79	0.92	-0.93	-1.86
PMP4/6-31G*	-6.85	-2.48	-0.77	1.19	0.38	-1.09	-1.88

^a Energies in kcal/mol.**Table VIII.** Relative Energies of OH Addition to C₂H₄ with 3-21G Basis Sets^a

level	R =						
	1.50	1.70	2.00	2.20	2.40	2.70	2.75
Without Spin Projection							
UHF/3-21G	-10.25	0.02	4.56	1.80	-0.25	-1.63	-1.73
MP2/3-21G	-18.96	-9.60	7.05	5.41	2.85	0.17	-0.13
MP3/3-21G	-15.96	-6.70	7.51	5.41	2.70	0.04	-0.25
MP4/3-21G	-16.59	-8.40	5.90	4.27	1.88	-0.41	-0.65
After Annihilation of the Largest Spin Contaminant							
PUHF/3-21G	-11.70	-5.59	-6.59	-6.04	-5.47	-4.22	-3.99
PMP2/3-21G	-19.97	-14.14	-2.72	-1.58	-1.84	-2.20	-2.22
PMP3/3-21G	-16.61	-9.83	-0.11	-0.07	-0.96	-1.79	-1.86
PMP4/3-21G	-17.23	-11.53	-1.71	-1.19	-1.78	-2.24	-2.25

^a Energies in kcal/mol.**Table IX.** Relative Energies of OH Addition to C₂H₄ with 6-31G* Basis Sets^a

level	R =						
	1.80	1.90	2.00	2.10	2.20	2.40	2.70
Without Spin Projection							
UHF/6-31G*	6.53	8.68	8.26	6.71	4.94	2.01	-0.36
MP2/6-31G*	-5.06	3.21	6.60	7.02	6.16	3.58	0.45
MP3/6-31G*	-1.45	5.32	7.76	7.62	6.41	3.53	0.37
MP4/6-31G*	-3.41	3.26	5.93	6.06	5.10	2.62	-0.08
After Annihilation of the Largest Spin Contaminant							
PUHF/6-31G*	-2.44	-3.03	-3.02	-2.97	-3.02	-3.15	-2.76
PMP2/6-31G*	-12.99	-7.33	-3.71	-1.96	-1.34	-1.46	-2.13
PMP3/6-31G*	-7.30	-2.82	-0.39	0.43	0.39	-0.55	-1.76
PMP4/6-31G*	-9.26	-4.88	-2.22	-1.12	-0.92	-1.46	-2.21

^a Energies in kcal/mol.

of a kcal/mol. Previous theoretical studies on similar complexes^{40,41} indicate that further increase in the basis set size will contribute less than 0.5 kcal/mol. Correlation corrections appear to change the binding energy by only 0.1–0.5 kcal/mol. Effects due to spin contamination are negligible.

Barrier Heights. The experimental activation energies are 1.3 kcal/mol for OH + C₂H₂ and -0.9 kcal/mol for OH + C₂H₄.^{9,10,13–16} Since the barrier heights are correlated with the activation energies, this would indicate that the experimental classical barrier heights are approximately 1 ± 2 and 0 ± 1 kcal/mol, respectively.⁴⁹ The theoretical data are collected in Tables VI–IX and plotted in Figures 4–7. At the UHF/3-21G level, the barrier heights are overestimated by 7 kcal/mol for OH + C₂H₂ and 5 kcal/mol for OH + C₂H₄. The addition of d orbitals to the basis set (3-21G to 6-31G*) increases the barrier heights by ca. 4 kcal/mol.

Because corrections for electron correlation and spin projection change and the position of the barriers, the barrier heights at the

(49) Truhlar, D. G., private communication.

Table X. Interpolated Barrier Heights^a

level	OH + C ₂ H ₂	OH + C ₂ H ₄
UHF/3-21G	8.3	4.6
MP2/3-21G	15.2	7.6
MP3/3-21G	12.4	7.8
MP4/3-21G	9.7	6.4
PUHF/3-21G	0.2	-4.5
PMP2/3-21G	2.6	-1.5
PMP3/3-21G	3.0	0.0
PMP4/3-21G	0.7	-1.2
UHF/6-31G*	12.0	8.7
MP2/6-31G*	16.3	7.0
MP3/6-31G*	14.5	8.0
MP4/6-31G*	11.2	6.1
PUHF/6-31G*	2.4	
PMP2/6-31G*	1.6	-1.2
PMP3/6-31G*	2.8	0.5
PMP4/6-31G*	1.6	-0.9
experiment	1.3 ± 0.1 ^b	-0.9 ± 0.3 ^c

^a In kcal/mol. ^b References 9 and 10. ^c References 13–16.

MPn and PMPn levels must be obtained by interpolation. This can be done graphically with Figures 4–7 or numerically with the data in Tables VI–IX. The interpolated barrier heights are summarized in Table X. For OH + C₂H₂ correlation corrections make the agreement with experiment worse, raising the barrier by as much as 7 kcal/mol. For OH + C₂H₄, the barriers at the correlated levels occur considerably earlier along the reaction path and are somewhat narrower. Correlation corrections with the 6-31G* basis improve the barrier heights only slightly.

Spin projection results in significant changes in the shapes of the potential energy curves and a remarkable improvement in the barrier heights: a decrease of 8–15 kcal/mol for OH + C₂H₂ and 7–9 kcal/mol for OH + C₂H₄. As noted before, the PUHF barriers and potential energy curves are generally not realistic.^{28,29,50} However, the shapes of the PMP2, PMP3, and PMP4 curves are quite reasonable and yield very similar barriers.⁵¹ The barriers are much lower and wider than at the unprojected levels and occur earlier along the reaction path. At the PMP4/6-31G* level, the barriers are 1.6 kcal/mol for OH + C₂H₂ and -0.9 kcal/mol for OH + C₂H₄. If transition-state theory is used to estimate the activation energies, the calculated activation energies at 298 K are 2.3 kcal/mol for OH + C₂H₂ and 0.1 kcal/mol for OH + C₂H₄ at the PMP4/6-31G* level,⁵² in good agreement with the experimental activation energies, 1.3 ± 0.1 and -0.9 ± 0.3 kcal/mol, respectively. Larger basis sets, more extensive treatment of correlation corrections, full optimization of the transition structures at a spin projected level, and the use of variation transition-state theory can be expected to further improve the agreement with experiment.

Conclusions

Barrier heights for OH + C₂H₂ and OH + C₂H₄ are too high at the UHF and UMPn levels because of spin contamination. Annihilation of the largest spin contaminant shifts the barriers toward the reactants by 0.1–0.3 Å and lowers the barriers by 7–15 kcal/mol. When corrections for spin projection are combined with electron correlation corrections, good agreement with the experimental activation energies is obtained. In addition to transition states, loose clusters were found for both OH + C₂H₂ and OH + C₂H₄.

Acknowledgment. This work was supported by a grant from the National Science Foundation (Grant No. CHE-83-12505).

(50) Mayer, I. *Adv. Quantum Chem.* 1980, 12, 189.

(51) Annihilation of the higher spin contaminants caused no significant changes in the positions, shapes, and heights of the barriers at the PMPn/3-21G level.

(52) Frequencies and thermal corrections were calculated at the HF/3-21G level with the HF/3-21G optimized structure at R = 2.2 Å (i.e., near the positions of the maxima in the PMP4 curves). The OH torsion modes have been treated as free rotors in the computation of the activation energies.